

Supercurrent Flow in NJL₂₊₁ at High Baryon Density

Simon Hands and Avtar Singh Sehra

*Department of Physics, University of Wales Swansea,
Singleton Park, Swansea SA2 8PP, U.K.*

Abstract: We present results of numerical simulations of the 2+1d Nambu – Jona-Lasinio model with non-zero baryon chemical potential μ and spatially-varying complex diquark source strength j . By choosing $\arg(j)$ to vary smoothly through 2π across the spatial extent of the lattice, a baryon number current is induced which in the high density phase remains non-vanishing as $|j| \rightarrow 0$; we are hence able to extract a quantity characteristic of a superfluid known as the *helicity modulus*. We also study supercurrent flow at non-zero temperature and estimate the critical temperature at which the normal phase is restored, which is consistent with the conventional picture for thin-film superfluids in which the transition is viewed in terms of vortex – anti-vortex unbinding.

1 Introduction

There are unfortunately rather few quantum field theories amenable to study using lattice Monte Carlo techniques in the presence of a non-zero chemical potential μ , or more specifically with $\mu/T \gg 1$. Many important theories, including QCD, cannot be studied because their path integral measure with $\mu \neq 0$ is not real on analytic continuation to Euclidean metric, making Monte Carlo importance sampling inoperative. Of those with positive definite measure, the Nambu Jona-Lasinio (NJL) model with $N_f = 2$ quark flavors [1] is one of the most interesting. At $\mu = 0$ the theory exhibits dynamical chiral symmetry breaking, with the generation of a constituent quark mass scale Σ much larger than the bare mass m . For $\mu > \mu_c \approx \Sigma$, chiral symmetry is restored, and the ground state is a degenerate fermi system with $\mu = E_F \simeq k_F$ [2]. In $d+1$ dimensions the baryon density in this case is $n_B = 4N_f \mu^d \theta(\mu - \mu_c) / (4\pi)^{\frac{d}{2}} d\Gamma(\frac{d}{2})$.

The precise nature of the ground state at high density depends on d . For the realistic case $d = 3$, lattice simulations suggest that condensation of diquark pairs at the Fermi surface takes place leading to spontaneous breakdown of the $U(1)_B$ baryon number symmetry [3]. An energy gap $\Delta > 0$ to excite fermionic quasiparticles develops; for phenomenologically-motivated lattice parameters the simulations predict $\Delta/\Sigma \simeq 0.15$, in good agreement with self-consistent model calculations of the gap in superconducting quark matter [4]. In this case the NJL model appears to behave as an orthodox BCS superfluid; there is long-range ordering of the ground state signalled by the non-vanishing

condensate $\langle qq \rangle \neq 0$, and a dynamically-generated mass scale Δ . Since a $U(1)_B$ symmetry has been spontaneously broken, there is a massless scalar qq bound state in the spectrum, which is associated with long-range interactions between vortex excitations in the superfluid, and with a collective propagating mode for $T > 0$ known as second sound.

However, both for obvious numerical convenience, and for a more formal reason, namely the existence of an interacting continuum limit, lattice simulations were first applied to the NJL model with $\mu \neq 0$ in 2+1 dimensions [5, 6, 7]. In this case the physics appears radically different. Whilst there is evidence for long-range coherence of diquark correlation functions [5], there is no long-range order, and apparently no gap. Rather, the condensate vanishes non-analytically with the diquark source strength, $\langle qq \rangle \propto j^\alpha$, with $0 < \alpha(\mu) < 1$ [6]. The results were interpreted in terms of a critical phase for all $\mu > \mu_c$, in which the diquark correlator decays algebraically, $\langle qq(0)qq(\vec{r}) \rangle \propto r^{-\eta}$ [7]. Since all simulations are performed on finite systems, and therefore necessarily at $T > 0$, the absence of long-range order is consistent with the Coleman-Mermin-Wagner theorem for $2d$ systems [8]. The situation is analogous to the low- T phase of the $2d$ X-Y model, one of whose physical applications is the description of superfluidity in thin films [9].

The defining property of a superfluid is that the flux density of conserved charge, or supercurrent $\vec{J}$, is related to the spatial variation of the phase angle θ of the $U(1)$ -valued order parameter field (in this case $\langle qq \rangle$) via

$$\vec{J} = \Upsilon \vec{\nabla} \theta. \quad (1)$$

The constant of proportionality Υ is known as the *helicity modulus*. For a textbook non-relativistic superfluid such as ^4He it is given by

$$\Upsilon = \frac{\hbar}{M} n_s \quad (2)$$

where M is the mass of the helium atom and n_s is a parameter called the *superfluid density*, which need not coincide with the charge density of the condensate. For a relativistic system Υ is best thought of as a phenomenological parameter in its own right, rather like f_π^2 in $(d+1)$ -dimensional chiral model [10]. One way of understanding superfluidity in the absence of long-range order in a $2d$ system is to observe that the only way to change the quantised circulation $\oint \vec{J} \cdot d\vec{l}$ around one direction of a finite torus is to excite a vortex – anti-vortex pair, and transport one member of the pair around the other direction of the torus before re-annihilation. The energy required to do this scales as $\ln L$ where L is the size of the system [9]: hence in the thermodynamic limit circulation patterns are topologically stable.

This Letter will present further support for this scenario in NJL_{2+1} by extracting Υ via a calculation of the induced baryon number current $\vec{J} = \langle \bar{\psi} \vec{\gamma} \psi \rangle$ in response to a spatially varying diquark source. As well as providing direct verification of superfluid behaviour in a fermionic model, we will also study the behaviour of $\vec{J}$ as temperature T is increased, and find the transition to “normal” behaviour at a critical T_c consistent with analytic expectations.

2 Method

The lattice NJL model studied is identical to that of [5, 7]:

$$S_{NJL} = \sum_x \bar{\chi}_x M[\Phi]_{xy} \chi_y + j \chi_x^{tr} \tau_2 \chi_x + \bar{j} \bar{\chi}_x \tau_2 \bar{\chi}_x^{tr} + \frac{1}{g^2} \sum_{\tilde{x}} \text{tr} \Phi_{\tilde{x}}^\dagger \Phi_{\tilde{x}}, \quad (3)$$

where $\chi, \bar{\chi}$ are isodoublet staggered lattice fermion fields, $\Phi = \sigma \mathbb{1} + i \vec{\pi} \cdot \vec{\tau}$ is an auxiliary bosonic field defined on the dual lattice sites $\tilde{x}$, and the matrix M is

$$M_{xy}^{pq} = \delta^{pq} \sum_{\nu=0,1,2} \frac{\eta_{\nu x}}{2} [e^{\mu \delta_{\nu 0}} \delta_{y, x+\hat{\nu}} - e^{-\mu \delta_{\nu 0}} \delta_{y, x-\hat{\nu}}] + \delta_{xy} \left\{ m \delta^{pq} + \frac{1}{8} \sum_{\langle \tilde{x}, x \rangle} [\sigma_{\tilde{x}} \delta^{pq} + i \varepsilon_x \vec{\pi}_{\tilde{x}} \cdot \vec{\tau}^{pq}] \right\}. \quad (4)$$

Here $\langle \tilde{x}, x \rangle$ denotes the set of 8 dual sites $\tilde{x}$ surrounding x , $\eta_{\mu x} = (-1)^{x_0 + \dots + x_{\mu-1}}$ is the Kawamoto-Smit phase required for a Lorentz covariant continuum limit, and $\varepsilon_x = (-1)^{x_0 + x_1 + x_2}$. A full description of the symmetries of (3) and the numerical simulation method is given in [7]. The only novelty in the current study is that the diquark source strengths $j, \bar{j}$ are now specified to be spatially varying, or “twisted”:

$$j = j_0 \exp(i\theta_{\tilde{x}}); \quad \bar{j} = j_0 \exp(-i\theta_{\tilde{x}}) \quad (5)$$

with j_0 a real constant. To ensure homogeneity and single-valuedness on an $L_s^2 \times L_t$ lattice we demand

$$\theta = \frac{2\pi}{L_s} (n_1 x_1 + n_2 x_2) \Rightarrow \vec{\nabla} \theta = \frac{2\pi}{L_s} (n_1, n_2). \quad (6)$$

A constant supercurrent of the form (1) is therefore specified by a pair of integers (n_1, n_2) . It remains to define the conserved baryon number current J_ν :

$$J_{\nu x} = \frac{1}{2} \langle e^{\mu \delta_{\nu 0}} \bar{\chi}_x \chi_{x+\hat{\nu}} + e^{-\mu \delta_{\nu 0}} \bar{\chi}_x \chi_{x-\hat{\nu}} \rangle. \quad (7)$$

The timelike component of (7) is none other than the baryon charge density n_B reported in [5, 7]. Here we shall use the same stochastic technique to estimate the quantum expectation value of the spacelike components $\vec{J}(j, \mu) = (L_s^2 L_t)^{-1} \sum_x \vec{J}_x(j, \mu)$ to demonstrate behaviour of the form (1). The strategy will be to compute $\vec{J}$ for fixed (n_1, n_2) for a range of j_0 and extrapolate $j_0 \rightarrow 0$. Behaviour consistent with (1) in this limit is deemed to be superfluid.

We used the same simulation parameters as [5, 7], namely $g^2 = 2.0a$, $ma = 0.01$, which at $\mu = 0$ yields a dynamically-generated constituent mass, which in effect sets the scale, of $\Sigma a = 0.71$. As μ is raised, there is a sharp transition from a chirally broken vacuum with $\langle \bar{\chi} \chi \rangle \simeq \frac{2}{g^2} \Sigma$, $n_B \simeq 0$ to a chirally restored phase with $n_B > 0$ at $\mu_c a \simeq 0.65$. Studies of the fermion dispersion relation in the phase $\mu > \mu_c$ are consistent with a sharp Fermi surface with $k_F \lesssim \mu$ and vanishing gap $\Delta \simeq 0$ [6, 7].

3 Results

3.1 $T = 0$




Figure 1: J_2 vs. j_0 on a $16^2 \times L_t$ lattice for two different values of μ .

In Fig. 1 we plot J_2 (strictly its imaginary part) as a function of j_0 for lattices of various temporal extent L_t at two representative values of μ : $\mu a = 0.2$ lies in the chirally-broken low density phase, and $\mu a = 0.8$ in the high density phase, where $n_B a^2 \simeq 0.25$ [7]. In all the plots shown here we have chosen $(n_1, n_2) = (0, 1)$ to minimise lattice artifacts, and from now on we set the lattice spacing $a = 1$.

The contrast between the two phases is quite dramatic. For $\mu = 0.2$ J_2 appears to vary approximately quadratically with j_0 , and extrapolate to zero as $j_0 \rightarrow 0$. There is no significant effect as $L_t \rightarrow \infty$, or alternatively as $T \rightarrow 0$. At $\mu = 0.8$ the small- j_0 behaviour depends very sensitively on L_t ; as $T \rightarrow 0$ the data accumulate on a straight line which clearly extrapolates to a non-zero value as $j_0 \rightarrow 0$.

This behaviour is readily explained by writing the order parameter (diquark) field as $\phi = \phi_0 e^{i\theta}$, with ϕ_0 approximately constant. A natural effective Hamiltonian for long wavelength order parameter fluctuations at low temperature is then

$$H_{eff} = \frac{1}{2}(\vec{\nabla}\phi)^* \cdot (\vec{\nabla}\phi) \simeq \frac{\phi_0^2}{2}(\vec{\nabla}\theta)^2. \quad (8)$$

The corresponding Noether current is $\vec{J} = -\frac{i}{2}[\phi^* \vec{\nabla}\phi - (\vec{\nabla}\phi^*)\phi] \simeq \phi_0^2 \vec{\nabla}\theta$. For $\mu < \mu_c$,

it is natural to postulate ϕ proportional to j , leading to $J_2(j_0) \propto j_0^2$. For $\mu > \mu_c$, if we assume that $\lim_{j_0 \rightarrow 0} \phi_0 \neq 0$ we recover (1) with $\Upsilon = \phi_0^2$.




Figure 2: J_2 vs. j_0 on a $16^2 \times 64$ lattice for various μ .

With confidence that $L_t = 64$ suffices to determine the phase, in Fig. 2 we plot $J_2(j_0)$ for various μ . There is a sharp change between values of $\mu \leq 0.65$, which display the low density quasi-quadratic behaviour and smoothly extrapolate to zero as $j_0 \rightarrow 0$, and $\mu \geq 0.68$ which show a negative curvature characteristic of the high density phase. We thus determine the critical chemical potential for the onset of superfluidity $0.65 < \mu_c < 0.68$, in good agreement with the critical value for chiral symmetry restoration. Since as yet we have no systematic method of extrapolating to $j_0 \rightarrow 0$ for $\mu \gtrsim \mu_c$ to obtain an estimate for $J_2(\mu)$ as an “order parameter”, we can make no decisive statement about the nature of the transition, but note that the behaviour of $J_2(j_0)$ varies much more sharply across the transition than the diquark condensate $\langle qq_+(j) \rangle$, either in this model (Cf. Figure 2 of [7]), or even in NJL₃₊₁ (Cf. Figure 4 of [3]). This matches the sharp drop in the chiral order parameter $\langle \bar{\chi}\chi \rangle$ and corresponding rise in n_B at $\mu = \mu_c$ [5, 7], and is consistent with the analytic prediction of a strong first order transition in the large- N_f limit [11].

From now on we work exclusively at $\mu = 0.8$ in an attempt to understand the superfluid phase further. In Fig. 3 we plot $\Upsilon = J_2 L_s / 2\pi$ versus j_0 on $L_s^2 \times 64$ lattices. Just as for the $\langle qq_+(j) \rangle$ data [7], it turns out that the data are well-fitted by $\Upsilon(j_0) = A + B/L_s$, resulting in the $L_s \rightarrow \infty$ extrapolation shown in the plot. Recall that as well as genuine finite-size effects in this case there may also be some discretisation artifacts, since as L_s increases the gradient operator in (6) becomes better-approximated by the finite




Figure 3: Υ vs. j_0 on a $L_s^2 \times 64$ lattice for $\mu = 0.8$.

difference. Finally the $\Upsilon(j_0)$ data in the thermodynamic limit are extrapolated to $j_0 \rightarrow 0$ with a remarkably simple linear fit, resulting in $\Upsilon = 0.1413(14)$. We thus quote a result for the helicity modulus of $\Upsilon/\Sigma = 0.200(2)$ at $\mu a = 0.8$.

It is interesting to pause and ask what value might be expected for Υ in a conventional symmetry-breaking scenario. Let us define diquark operators $qq_{\pm} = \frac{1}{2}(\chi^{tr}\tau_2\chi \pm \bar{\chi}\tau_2\bar{\chi}^{tr})$ and source strengths $j_{\pm} = j \pm \bar{j}$, so that the diquark terms in the action (3) read $j_+qq_+ + j_-qq_-$. In the limit $j_- = 0$ the equation of motion for the current is then

$$\Delta_{\mu}^{-}J_{\mu} = 2j_+qq_-. \quad (9)$$

In the same limit the $U(1)_B$ -equivalent of the axial Ward identity reads

$$\langle qq_+ \rangle = j_+ \sum_x \langle qq_-(0)qq_-(x) \rangle = \frac{j_+}{M_-^2} |\langle 0|qq_-|- \rangle|^2. \quad (10)$$

where the second equality assumes that the correlation function is dominated by a pseudo-Goldstone pole of the form $(k^2 + M_-^2)^{-1}$, and $|- \rangle$ denotes a one-Goldstone state. We now introduce the $U(1)_B$ -equivalent form of the PCAC hypothesis:

$$\langle 0|\Delta_{\mu}^{-}J_{\mu}|- \rangle = \sqrt{\Upsilon}M_-^2 = 2j_+\langle 0|qq_-|- \rangle \quad (11)$$

where we have used the relation $\Upsilon = f_{\pi}^2$ derived in [10], and the second equality follows from (9). Combining (10) and (11) leads to the equivalent of the ‘‘Gell-Mann-Oakes-Renner’’ relation:

$$\Upsilon_{GMOR}M_-^2 = 8j\langle qq_+ \rangle. \quad (12)$$

This can be compared with numerical data for $\langle qq_+(j) \rangle$ and $M_-(j)$ in [7]. At $j = 0.3$, $M_- = 0.95$, $\langle qq_+ \rangle = 0.72$ yielding $\Upsilon_{GMOR} \approx 1.9$; at $j = 0.1$, $M_- = 0.4$, $\langle qq_+ \rangle = 0.52$ yielding $\Upsilon_{GMOR} \approx 2.6$. We conclude $\Upsilon \ll \Upsilon_{GMOR}$, consistent with our hypothesis that no symmetry breaking occurs, but that the dynamics are dominated by long-range phase fluctuations of the order parameter field, described by a strongly-interacting scalar diquark field rather than a weakly-interacting Goldstone mode.

3.2 $T > 0$

In this section for the first time we explore the superfluid phase at non-zero temperature. We expect a restoration to the normal phase at some critical T_c . In a comparable numerical study of NJL₃₊₁ which exhibits superfluidity via orthodox symmetry breaking [3], the value of T_c could be estimated from the zero temperature gap Δ via the BCS prediction $\Delta/T_c \simeq 1.76$. Since this implied that L_t had to exceed $35a$ for the system to be superfluid, an unambiguous extrapolation $j \rightarrow 0$ to permit a systematic study of $T > 0$ was not possible. In the current case we shall see that although the $j \rightarrow 0$ extrapolation still remains a problem, attaining $T < T_c$ is well within reach of the simulation.

First let us review a heuristic argument for the expected value of T_c , starting from the Hamiltonian H_{eff} (8) with $\phi_0^2 = \Upsilon$ [9]. The phase field $\theta(\vec{x})$ may be disrupted by topologically non-trivial vortex excitations of the form $\theta = q\psi$, $|\vec{\nabla}\theta| = q/r$, where q is integer and $\vec{x}$ is written (r, ψ) . The energy of a single vortex is thus

$$E \approx \frac{\Upsilon}{2} \int_a^{L_s} 2\pi r dr \left(\frac{q}{r}\right)^2 = \pi \Upsilon q^2 \ln\left(\frac{L_s}{a}\right). \quad (13)$$

Since a vortex can be located on any of L_s^2 lattice sites, the entropy

$$S = 2 \ln\left(\frac{L_s}{a}\right). \quad (14)$$

The free energy $F = E - TS$ thus changes sign for $q = 1$ vortices at a critical temperature

$$T_c = \frac{\pi}{2} \Upsilon. \quad (15)$$

The interpretation is that a phase transition separates a low- T superfluid phase in which vortices are confined to bound dipole pairs, and a high- T normal phase in which the vortex anti-vortex plasma screens the long-range interactions responsible for the divergent energy in (13). The relation (15) remains valid in a more sophisticated renormalisation group treatment, except that Υ is now T -dependent and should be replaced by its value $\Upsilon(T_c)$ exactly at the transition [12].

Combining our result for Υ with (15) yields an estimate $L_t \approx 4.5$ for the temporal lattice extent where a transition to the normal phase might be expected at $\mu = 0.8$. Fig. 4 shows $J_2(j_0)$ on $32^2 \times L_t$ lattices with L_t ranging from 64 all the way down to 2. At the extremes $L_t \geq 32$, $L_t \leq 4$ the data are reminiscent of those characterising respectively the




Figure 4: J_2 vs. j_0 on a $32^2 \times L_t$ lattice at $\mu = 0.8$ for various L_t .

high and low baryon density phases in Fig. 1. For intermediate temperatures, however, $J_2(j_0)$ shows positive curvature near the origin followed by negative curvature at larger j_0 , and once again the means of extrapolating $j_0 \rightarrow 0$ to determine whether superfluidity




Figure 5: J_2 vs. L_t on a $32^2 \times L_t$ lattice at $\mu = 0.8$ for fixed $j_0 = 0.025$.

persists is unclear.

In Fig. 5 we try a different tactic, plotting J_2 for every even $L_t \in [2, 64]$ for fixed $j_0 = 0.025$. A linear fit $J_0 = a_0 L_t + a_1$ through data with $L_t \leq 42$ seems quite reasonable, yielding $a_0 = 0.00212(25)$, $a_1 = -0.01537(9)$. If we identify the intercept on the L_t -axis with the transition, we deduce $L_{tc} = 7.25(95)$ and hence $\Upsilon/T_c = 1.02(13)$, to be compared with the theoretical value 0.637 from (15).

4 Summary

In this short study of the response of the system to a twisted diquark source forcing a baryon number current, we have provided direct evidence for the superfluid nature of the ground state of NJL₂₊₁ at high baryon density, and quantified it at one representative value of μ via the helicity modulus Υ . It should be stressed that the “physical” value $\Upsilon/\Sigma \simeq 0.2$ quoted is still to be extrapolated to the continuum limit. It is probably more important to note that the numerical value of Υ is an order of magnitude smaller than might be expected in an orthodox symmetry breaking scenario, and is consistent with the non-Goldstone, strongly self-interacting nature of the scalar diquark excitations above the ground state.

We also studied the response of the system to non-zero temperature. Whilst we were unable to extrapolate to the zero source limit in a controlled way, by studying fixed j_0 we were able to estimate a critical temperature T_c for breakdown of superfluidity of the same order of magnitude as, and only slightly smaller than, the Kosterlitz-Thouless prediction for a $2d$ system with U(1) global symmetry, which follows from characterising the superfluid/normal transition as arising from vortex pair unbinding. More refined simulations would be required to determine whether T_c actually has the KT value, or whether NJL₂₊₁, which in addition to the scalar diquark excitations contains massless fermion degrees of freedom, actually lies in a different universality class, as suggested by estimates of the critical exponent δ [7].

Acknowledgements

SJH was supported by a PPARC Senior Research Fellowship. We are grateful to Costas Strouthos for valuable insight in the early stages of the project.

References

- [1] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 345; *ibid* **124** (1961) 246.
- [2] S.P. Klevansky, Rev. Mod. Phys. **64** (1992) 649.
- [3] S.J. Hands and D.N. Walters, Phys. Rev. **D69** (2004) 076011.

- [4] K. Rajagopal and F. Wilczek, in *Handbook of QCD* (World Scientific, 2001), ch. 35, [arXiv:hep-ph/0011333](#); M.G. Alford, Ann. Rev. Nucl. Part. Sci. **51** (2001) 131.
- [5] S.J. Hands and S.E. Morrison, Phys. Rev. **D59** (1999) 116002.
- [6] S.J. Hands, B. Lucini and S.E. Morrison, Phys. Rev. Lett. **86** (2001) 753.
- [7] S.J. Hands, B. Lucini and S.E. Morrison, Phys. Rev. **D65** (2002) 036004.
- [8] N.D. Mermin and H. Wagner, Phys. Rev. Lett. **17** (1966) 1133;
S. Coleman, Comm. Math. Phys. **31** (1973) 259.
- [9] J.M. Kosterlitz and D.J. Thouless, J. Phys. **C6** (1973) 1181.
- [10] P. Hasenfratz and H. Leutwyler, Nucl. Phys. B **343**, 241 (1990).
- [11] K.G. Klimenko, Z. Phys. **C37** (1988) 457;
B. Rosenstein, B.J. Warr and S.H. Park, Phys. Rev. **D39** (1989) 3088.
- [12] D.R. Nelson and J.M. Kosterlitz, Phys. Rev. Lett. **39** (1977) 1201;
D.R. Nelson, in *Phase Transitions and Critical Phenomena*, Vol **7** (1983) p.1, eds.
C. Domb and J.L. Lebowitz (Academic Press, London).